
















Acetaminophen dosing directions for infants and children 11 years and younger

Acetaminophen	If possible, use weight to dose; otherwise use age.							
	6-11 lbs 0-3 mos	12-17 lbs 4-11 mos	18-23 lbs 12-23 mos	24-35 lbs 2-3 yrs	36-47 lbs 4-5 yrs	48-59 lbs 6-8 yrs	60-71 lbs 9-10 yrs	72-95 lbs 11 yrs
Infants' acetaminophen drops for example Infants' Tylenol® Oral Suspension 160 mg per 5 mL	Today's date _____  _____ mL _____ mL _____ mL Ask your doctor for the right dose (expires in 14 days)			 5 mL	—	—	—	—
Children's acetaminophen liquid for example Children's Tylenol® Oral Suspension 160 mg per 5 mL	← Ask a doctor →			 5 mL	 7.5 mL	 10 mL	 12.5 mL	 15 mL
Children's 80 mg acetaminophen tablets 80 mg per chewable tablet	← Ask a doctor →			 2 chewables	 3 chewables	 4 chewables	 5 chewables	 6 chewables
Children's or Junior 160 mg acetaminophen tablets 160 mg per chewable tablet	← Ask a doctor →			 2 chewables	 2½ chewables	 3 chewables		

Important instructions

You may repeat the dose every 4 hours while symptoms last

Do **NOT** give more than 5 doses in 24 hours

Do **NOT** use with any other product containing acetaminophen














- What you can do**
- Always read and follow the medicine label.
 - Find the dose for your child's **current weight**. If you are not sure of your child's weight, use age. As children grow, their doses change over time.
 - Use **only the dosing device that came with the medicine** (syringe or dosing cup).
 - Wait the **full amount of time between doses**. If you give another dose too soon, it could harm your child.

See [ibuprofen dosing directions](#) ►

Find tips, quizzes, and videos at [GetReliefResponsibly.com](https://www.getreliefresponsibly.com)

Get Relief Responsibly®

Ibuprofen dosing directions for infants and children 11 years and younger

Ibuprofen	If possible, use weight to dose; otherwise use age.							
	— <6 mos	12-17 lbs 6-11 mos	18-23 lbs 12-23 mos	24-35 lbs 2-3 yrs	36-47 lbs 4-5 yrs	48-59 lbs 6-8 yrs	60-71 lbs 9-10 yrs	72-95 lbs 11 yrs
Infants' ibuprofen concentrated drops <i>for example</i> Infants' Motrin® or Advil® Concentrated Drops 50 mg per 1.25 mL	Ask a doctor	 1.25 mL	 1.875 mL	—	—	—	—	—
Children's ibuprofen liquid <i>for example</i> Children's Motrin® or Advil® Oral Suspension 100 mg per 5 mL	Ask a doctor			 5 mL	 7.5 mL	 10 mL	 12.5 mL	 15 mL
Junior strength ibuprofen chewables <i>for example</i> Junior Strength Advil® Chewables 100 mg per chewable tablet	Ask a doctor					 2 chewables	 2½ chewables	 3 chewables
Junior strength ibuprofen tablets <i>for example</i> Junior Strength Advil® Tablets 100 mg per tablet	Ask a doctor					 2 tablets	 2 tablets	 3 tablets

Important instructions

You may repeat the dose every 6 to 8 hours while symptoms last

Do **NOT** give more than 4 doses in 24 hours

Talk to your child's doctor before giving any other product containing an NSAID*

*nonsteroidal anti-inflammatory drug

What you can do

- Always read and follow the medicine label.
- Find the dose for your child's **current weight**. If you are not sure of your child's weight, use age. As children grow, their doses change over time.
- Use **only the dosing device that came with the medicine** (syringe or dosing cup).
- Wait the **full amount of time between doses**. If you give another dose too soon, it could harm your child.

See acetaminophen dosing directions ►

Find tips, quizzes, and videos at [GetReliefResponsibly.com](https://www.getreliefresponsibly.com)

Get Relief Responsibly®